U.S. Environmental Protection Agency

Nutrient Criteria Technical Guidance Manual

Estuarine and Coastal Marine Waters

October 2001

Disclaimer

This manual provides technical guidance to States, Indian Tribes, and other authorized jurisdictions to establish water quality criteria and standards under the Clean Water Act (CWA), to protect aquatic life from acute and chronic effects of nutrient overenrichment. Under the CWA, States and Indian Tribes are to establish water quality criteria to protect designated uses. State and Indian Tribal decisionmakers retain the discretion to adopt approaches on a case-by-case basis that differ from this guidance when appropriate and scientifically defensible. Although this manual constitutes EPA's scientific recommendations regarding ambient concentrations of nutrients that protect resource quality and aquatic life, it does not substitute for the CWA or EPA's regulations; nor is it a regulation itself. Thus, it cannot impose legally binding requirements on EPA, States, Indian Tribes, or the regulated community, and might not apply to a particular situation or circumstance. EPA may change this guidance in the future.

Cover Photograph: Somewhere on the Chesapeake Bay. Supplied by David Flemer as a duplicate copy from the Chesapeake Biological Laboratory Photo Archives, University of Maryland; date unknown but earlier than 1972.

CONTENTS

Contributors	
Acknowledgments	
Foreword	
Executive Summary	xvii
Chapter 1. Introduction and Objectives	1-1
1.1 Backround	1-1
1.2 Definition of Estuaries and Coastal Systems	
1.3 Nature of the Nutrient Overenrichment Problem in Estuarine and Coastal Marin	
Scope and Magnitude of the Problem	1-5
1.4 The Nutrient Criteria Development Process	
Preliminary Steps	1-8
Strategy for Reducing Human-Based Eutrophication	1-10
Nutrient Criteria Development Process	1-11
Chapter 2. Scientific Basis for Estuarine and Coastal Waters Quantitative Nut	rient Criteria 2-1
2.1 Introduction	2-1
Purpose and Overview	2-1
Some Important Nutrient-Related Scientific Issues	
River-to-Ocean Continuum: Watershed/Nearshore Coastal Management Frame	
2.2 Controlling the Right Nutrients	
Overview	
Some Empirical Evidence for N Limitation of Net Primary Production	
Some Threshold Responses to Nitrogen Overenrichment	
Effects of Physical Forcing on Net Primary Production	
Other Physical Factors	
2.3 Nutrient Loads and Concentrations: Interpretation of Effects	
Conceptual Framework	
Examples	
2.4 Physical-Chemical Processes and Dissolved Oxygen Deficiency	
2.5 Nutrient Overenrichment Effects and Important Biological Resources	
Benthic Vascular Plant Responses to Nutrients	
Other Examples of Important Biotic Effects of Nutrient Overenrichment	
2.6 Concluding Statement on Nitrogen and Phosphorus Controls	2-32
Chapter 3. Classification of Estuarine and Coastal Waters	
3.1 Introduction	
Purpose and Background	3-1
Defining the Resource of Concern	
3.2 Major Factors Influencing Estuarine Susceptibility to Nutrient Overenrichment	
Dilution	3-3
Water Residence Time	3-3
Stratification	3-3

3.3 Examples of Coastal Classification	3-4
Geomorphic Classification	3-4
Man-Made Estuaries	3-6
Physical/Hydrodynamic Factor–Based Classifications	3-6
Other Considerations	3-9
Summary	3-11
3.4 Coastal Waters Seaward of Estuaries	3-11
Geomorphic Classification	3-12
Nongeomorphic Classification	
Chapter 4. Variables and Measurement Methods To Assess and Monitor	
Estuarine/Marine Eutrophic Conditions	4-1
4.1 Introduction	
4.2 Causal and Response Indicator Variables	
Nutrients as Causal Variables	
Response Variables	4-6
Measures of Water Clarity	4-7
Dissolved Oxygen	4-8
Benthic Macroinfauna	4-9
4.3 Field Sampling and Laboratory Analytical Methods	
Field Sampling Methods	
Laboratory Analytical Methods	4-13
Water Column Nutrients	4-13
Sediment Analyses	4-15
Determination of Primary Productivity	4-16
Phytoplankton Species Composition	4-16
Macrobenthos, Macroalgae, and Seagrasses and SAV	4-17
	_
Chapter 5. Databases, Sampling Design, and Data Analysis	5-1
5.1 Introduction	5 _1
5.2 Developing Regional and National Databases for Estuaries and Coastal Waters	
Data Sources	
EPA Water Quality Data	
National Oceanographic and Atmospheric Administration (NOAA)	
· · · · · · · · · · · · · · · · · · ·	
Rivers and Streams Water Quality Data	
USGS San Francisco Bay Program	
State/Tribal Monitoring Programs	
Sanitation Districts	
Academic and Literature Sources	
Volunteer Monitoring Programs	
Quality of Historical Data	
Location Data	
Variables and Analytical Methods	
Laboratory Quality Control	
Data Collecting Agencies	
Time Period	5-10

	Index Period	. 5-10
	Representativeness	. 5-10
	Gathering New Data	. 5-10
5.3	Sampling Design	
	Sampling Protocol	. 5-11
	Sampling Technique	
	Initial Considerations	. 5-12
	Specifying the Population and Sample Unit	. 5-13
	Specifying the Reporting Unit	. 5-14
	Sources of Variability	
	Alternative Sampling Designs	
	Monitoring Programs	
	Citizen Monitoring Programs	
5.4	Quality Assurance/Quality Control	
	Representativeness	
	Completeness	
	Comparability	
	Accuracy	
	Variability	
5.5	Statistical Analyses	
	Data Reduction	
	Frequency Distributions	
	Correlation and Regression Analyses	
	Tests of Significance	
	apter 6. Determining the Reference Condition	
	Introduction and Definition	
	Significance of Reference Conditions	
	Paucity of Similar Estuarine and Coastal Marine Ecosystems	
6.4	Approaches for Establishing Reference Conditions	
	In Situ Observations as the Basis for Estuarine Reference Condition	
	Areal Load Approach to Identification of Reference Condition	. 6-11
Cha	apter 7. Nutrient and Algal Criteria Development	7-1
	Introduction	
	Role of Regional Technical Assistance Groups	
	Classification	
7.4	Descriptive Background Information	
	Estuarine Watershed Characterization	
	Within Estuarine System Characterization	
7.5	Elements of Nutrient Criteria	
	D C 11/2	
	Reference Condition	
	Historical Information	7-5
	Historical Information	7-5
	Historical Information	7-5 7-5

• •	etical Examples of Nutrient Criteria Development Deliberations	
	ioio	
7.7 Evaluati	ion of Proposed Criteria	7-8
	nce for Interpreting and Applying Criteria	
Do the	Criteria Protect Designated Uses?	7-9
	ation Goals	
Sampl	ing for Comparison to Criteria	7-11
7.8 Nutrient	t Criteria Interpretation Procedures	7-12
Decision	onmaking Protocol	7-12
Multiv	ariable Enrichment Index	7-12
	ency and Duration	
	Modifications	
	State, or Tribe Responsibility under the Clean Water Act	
7.11 Impler	mentation of Nutrient Criteria into Water Quality Standards	7-14
Chapter 8.	Using Nutrient Criteria to Protect Water Quality	8-1
0.1 Managir	n a Daint Causa Dallution	0 1
_	ng Point Source Pollution	
	lean Water Act and Water Quality Standards	
	ting Designated Uses	
	aining Existing Water Quality	
	al Policies	
	S Permits	
	o the Future Pollutant Trading	
	ing Nonpoint Source Politifier	
	s to Control Nonpoint Source Pollution	
	al Estuary Program	
	pheric Deposition	
	l Nonpoint Pollution Control Programs	
	Bill Conservation Provisions	
	hensive Procedure for Nutrient Management	
	Status Identification	
	Background Investigation	
	Data Gathering and Diagnostic Monitoring	
	Source Identification	
	: Management Practices for Nutrient Control	
_	Detailed Management Plan Development	
	: Implementation and Communication	
	Evaluation Monitoring and Periodic Review	
	: Completion and Evaluation	
	0: Continued Monitoring of the System	
	es	
0.4 Resourc		6-20
Chapter 9.	Use of Models in Nutrient Criteria Development	9-1
9.1 Introduc	ction	9-1
	Empirical Models in Nutrient Criteria Development	

Use of M	Iathematical Models in Nutrient Criteria Development	. 9-3
9.2 Model Ide	entification and Selection	. 9-3
Model Id	lentification	. 9-4
Model Se	election	. 9-8
9.3 Model Cla	assification	. 9-9
Level I N	Models	9-10
Level II	Models	9-13
Level III	Models	9-14
Level IV	Models	9-15
Summary	y of Model Capabilities	9-17
	odels for Nutrient Investigation	
	alibration and Validation	
9.5 Manageme	ent Applications	9-22
Load-Re	sponse Analysis	9-22
Acceptal	ole Nutrient Loads	9-23
Case Stu	dy Example	9-24
References		R-1
Appendixes		
Appendix A:	Conditions for Bloom Development: Interplay among Biogeochemical, Biological,	
11	and Physical Processes	A-1
Appendix B:	Additional Information on the Role of Temperature and Light on Estuarine	
11	and Coastal Marine Phytoplankton	B-1
Appendix C:	Additional Information on Flushing in Estuaries	
Appendix D:	NOAA Scheme for Determining Estuarine Susceptibility	
Appendix E:	Comparative Systems Empirical Modeling Approach: the Empirical Regression	
	Method to Determine Nutrient Load-Ecological Response Relationships	
	for Estuarine and Coastal Waters	
Appendix F:	Selected Theoretical Approaches to Classification of Estuaries and Coastal Waters	. F-1
Appendix G:	Examples of Nutrient Concentration Ranges and Related Hydrographic Data	
	for Selected Estuaries and Coastal Waters in the Contiguous States	
	of the United States	G-1
Appendix H:	Preliminary Statement of Proposed Near Coastal Marine Nutrient Sampling and	
	Reference Condition Development Procedure	H-1
Cose Studies		
Case Studies		
San Francisco	Bay Program: Managing Coastal Resources of the U.S	CS-1
	ound - Hypoxia	
	Narragansett Bay	
-	ase Study	
	sapeake Bay Water Quality	
_	from Washington State	
A I dispective	nom washington state	J-4Z

Figures

Figure 1-1a.	Draft aggregation of Level III ecoregions for the National Nutrient Strategy illustrating
	those areas most related to coastal and estuarine nutrient criteria development 1-3
Figure 1-1b.	Coastal provinces
Figure 1-2.	The eutrophication process
Figure 1-3.	Expanded nutrient enrichment model Source: Bricker et al. 1999 1-7
Figure 1-4.	Elements of nutrient criteria development and their relationships in the process 1-9
Figure 1-5.	Derivation of the reference condition and the National Nutrient Criteria Program
U	using TP, TN, and Chlorophyll <i>a</i> as example variables
Figure 1-6.	Flowchart of the nutrient criteria development process
Figure 2-1.	Idealized scheme defining the coastal ocean and the coastal zone2-2
Figure 2-2.	Schematic representation of contemporary (Phase II) conceptual model of coastal
118010 = 21	eutrophication
Figure 2-3.	Salinity zones
Figure 2-4.	Schematic illustrating the central role of phytoplankton as agents of biogeochemical
Tigure 2 4.	change in shallow coastal ecosystems
Figure 2-5.	Transport of nutrients to Laholm Bay, Sweden
Figure 2-6.	Summary of nitrogen:phosphorus ratios in 28 sample estuarine ecosystems 2-14
Figure 2-7.	Factors that determine whether nitrogen or phosphorus is more limiting in
rigule 2-7.	aquatic ecosystems
Eigung 2.9	
Figure 2-8.	Cartoon diagrams of three physical forcings that operate at the interface between SCEs
F: 2 0	and the coastal ocean (tides), watershed (river inflow), and atmosphere (wind) 2-19
Figure 2-9.	Simple schematic diagram showing the influences of river flow on ecosystem
E: 0.10	stocks and processes examined in this study
Figure 2-10.	Scatter diagram showing the relationship between the rate of decline in
	dissolved-oxygen concentrations in deep water and average deposition rates
F: 0.11	of total chlorophyll <i>a</i> during the spring-bloom period
Figure 2-11	Schematic diagram of coastal plain estuary types, indicating direction and
a-d.	degree of mixing
Figure 2-12.	Net transports in estuaries resulting from estuarine flows and mixing 2-23
Figure 2-13.	Net movement of a particle in each layer of a two-layered flow system 2-23
Figure 2-14.	The fraction of landside nitrogen input exported from 11 North American and
	European estuaries versus freshwater residence time (linear time scale) 2-25
Figure 2-15.	Scatter plots of water column averaged chlorophyll <i>a</i> at a mesohaline station versus
	several different functions of total nitrogen loading rate measured at the fall line of
	the Potomac River estuary
Figure 2-16a.	Primary production by phytoplankton as a function of the estimated annual input of
	dissolved inorganic nitrogen per unit volume of a wide range of marine ecosystems 2-30
Figure 2-16b.	Primary production by phytoplankton as a function of the annual input of dissolved
	inorganic nitrogen per unit area of a wide range of marine ecosystems 2-30
Figure 2-16c.	Fisheries yield per unit area as a function of primary production in a wide range of
	estuarine and marine systems
Figure 2-17.	Comparative evaluation of fishery response to nutrients
Figure 3-1.	Idealized micronutrient-salinity relations showing concentration and mixing of
-	nutrient-rich river water with nutrient-poor seawater
Figure 3-2a.	Relationship between the mean annual loadings of dissolved inorganic nitrogen
C	and the mean annual concentration of chlorophyll <i>a</i> in microtidal and macrotidal
	estuaries

Figure 3-2b.	Relationship between the mean annual concentrations of dissolved inorganic	
	nitrogen and chlorophyll a in microtidal and macrotidal estuaries	. 3-12
Figure 6-1.	Environmental quality scale representing reference conditions and potential nutrient criteria relative to designated uses	
Figure 6.2	Hypothetical frequency distribution of nutrient-related variables showing quantities for	
Figure 6-2.	reference or high-quality data and mixed data	
Figure 6-3.	Hypothetical example of load/concentration response of estuarine biota to increased	0-7
riguic 0-3.	enrichment	6-0
Figure 6-4.	An illustration of the comparison of past and present nutrient data to establish	0 /
Tiguic 0 4.	a reference condition for intensively degraded estuaries.	6-10
Figure 6-5.	Areal load estimate approach to nutrient reference condition determination	
Figure 7-1.	Generalized progression and relationship of the elements of a nutrient criterion	
Figure 7-2.	Hypothetical illustration of developing a TN criterion in an estuary	
Figure 8-1.	Components of water quality standards	
Figure 8-2.	"Threefold framework" of evaluation.	
Figure 9-1.	Eutrophication model framework	
Figure 9-2.	Use of models in load-response analysis	
Figure 9-3.	Use of models in determining allowable loads	
Figure 9-4.	Shipps Creek site map and salinity monitoring locations	
Figure 9-5.	Model results for existing conditions	
Figure 9-6.	Model results for 50% reduction in WWTP load	
	Tables	
Table 2-1.	Categorization of the world's continental shelves based on location, major river,	
	and primary productivity	2-8
Table 2-2.	Estuaries exhibiting seasonal shifts in nutrient limitation with spring P limitation	
m 11 00	and summer N limitation	
Table 2-3.	DO, nutrient loading, and other characteristics for selected coastal areas and a MERL	
T 11 2 1	mesocosm enrichment experiment	
Table 3-1.	General drowned river valley estuarine characteristics	3-5
Table 3-2.	Classification of coastal systems based on relative importance of river flow, tides,	2.0
Table 4-1.	and waves to mixing	3->
1 able 4-1.	coastal and marine environments	1 10
Table 6-1.	Summary of estuarine and coastal nutrient reference condition determinations	
Table 6-1.	Requisite assumptions for establishing watershed-based reference conditions	
Table 7-1.	Example of an enrichment index using the middle portion of a hypothetical estuary	
Table 8-1.	States for which the nonpoint source agency is not the water quality agency	
Table 8-2.	States and territories with coastal nonpoint pollution control programs	
Table 9-1.	Basic model features	
Table 9-2.	Key features of selected models	
Table 9-3.	Calculation spreadsheet for Shipps Creek estuary	
Table 9-4.	Chesapeake watershed nitrogen deposition under varying management schemes for	. , 20
	emissions of nitrogen atmospheric depositions precursors	. 9-29
Table 9-5.	Water quality state variables used in CBEMP	
	•	

CONTRIBUTORS

Jessica Barrera (Hispanic Association of Colleges and Universities/University of Miami)

Robert Cantilli (U.S. Environmental Protection Agency)

Ifeyinwa Davis (U.S. Environmental Protection Agency)*

Edward Dettmann (U.S. Environmental Protection Agency)

Jen Fisher (University of Georgia)

David Flemer (U.S. Environmental Protection Agency)*

Thomas Gardner (U.S. Environmental Protection Agency)*

George Gibson (U.S. Environmental Protection Agency)*

Debbi Hart (U.S. Environmental Protection Agency)

James Latimer (U.S. Environmental Protection Agency)

Scott Libby (Battelle)*

Greg Smith (GLEC, Inc.)

CarolAnn Siciliano (U.S. Environmental Protection Agency)*

Jack Word (MEC Analytical Systems)*

WORK GROUP MEMBERS

Ifeyinwa Davis (U.S. Environmental Protection Agency)

David Flemer (U.S. Environmental Protection Agency)

John Fox (U.S. Environmental Protection Agency)

George Gibson (U.S. Environmental Protection Agency)

Debbi Hart (U.S. Environmental Protection Agency)

Suzanne Bricker (National Oceanic and Atmospheric Administration)

Dorothy Leonard (National Oceanic and Atmospheric Administration)

Scott Libby (Battelle)

Greg Smith (GLEC, Inc.)

Jack Word (MEC Analytical Systems)

^{*}Denotes primary authors

ACKNOWLEDGMENTS

The authors wish to thank the following peer reviewers for their assistance in the preparation of this manual: Don Boesch (University of Maryland) and Hans Paerl (Institute of Marine Sciences, University of North Carolina), Matthew Liebman (U.S. EPA Region I), Michael Bira (U.S. EPA Region VI), and Kenneth Teague (U.S. EPA Region VI). Walter Nelson and Peter Eldridge (U.S. EPA Office of Research and Development) provided comments on an early draft of the manuscript, and Thomas Brosnan (National Oceanic and Atmospheric Administration, Damage Assessment Center), Michael Kemp (Horn Point Laboratory, CES, University of Maryland), Jonathan Pennock (Dauphin Island Laboratory, University of South Alabama), Hassan Mirsajadi (Delaware Department of Natural Resources and Environmental Control), and David Tomasko (State of Florida Southwest Florida Water Management District) provided formal peer review comments on a final working draft. Additional comments on the working draft were provided by Suesan Saucerman (U.S. EPA Region IX); John (Jack) Kelly, James Latimer, and Edward Dettmann (U.S. EPA ORD); Lewis Linker (U.S. EPA Chesapeake Bay Program); Laura Gabanski (U.S. EPA Office of Wetlands, Oceans, and Watersheds); Joel Salter (Office of Wastewater Management); Mimi Dannel and Marjorie Wellman (U.S. EPA Office Science and Technology); CarolAnn Siciliano (U.S. EPA Office of General Council); and Cynthia Moncreiff (University of Southern Mississippi). Treda Smith (U.S. EPA Office of Water) assisted in compiling references. Edits and suggestions made by the peer review panel were incorporated into the final version of the manual. State agencies and private interest groups also offered comments and they were addressed where possible in this manual. We appreciate the work of Joanna Taylor, The CDM Group, Inc., who patiently and graciously made repeated format changes to this manuscript as it evolved over the several months. The authors of the case studies are thanked for their contributions.

Estuaries and coastal waters are a diverse suite of ecosystems, and differences in methods and approaches are to be expected. This and subsequent manuals are not intended to be singular, one-time publications. As experience accumulates, future editions will be prepared to reflect new understanding.

FOREWORD

This manual is intended for State/Tribal and Federal agency personnel actively engaged in water resource management data collection, assessment, planning, and project implementation. Consequently, it incorporates both a scientific rationale and enough of the "nuts and bolts" of nutrient criteria development and management to help both initiates and those experienced in water resource management.

These nutrient criteria development and management efforts are directed at anthropogenic sources. Inherent "natural" background levels are not and should not be subject to management. Our responsibility is to abate human-caused eutrophication in estuaries and coastal or "near coastal" (out to about 20 nautical miles) marine waters.

To distinguish between natural background enrichment and human impacts, it is necessary to identify localities that experience minimal human influence. Ambient nutrient measurements at these sites may then be compared to similar sites that do experience human influences. The difference in nutrient measurements is the difference between a reference site and a test site. A reference condition is a collection of measurements from several reference sites that incorporates a central tendency statistic.

Because of differences in geologic parent material, climate, and geography, reference conditions are different from one region to another. Similarly, waterbodies, especially estuaries, often respond differently to nutrient inputs. Lakes and reservoirs are different from streams and rivers, and estuaries and coastal marine waters have characteristics different from both. Criteria have to be designed for particular waterbody types and the regions in which they lie.

The primary variables of concern in criteria development are two causal enrichment variables: total phosphorus (TP) and total nitrogen (TN). These nutrients are essential to algal and plant production and are the base of the food chain that supports all other life in the system. Also, two initial response variables usually are the first indicators of biological growth reaction to enrichment. One is chlorophyll *a*, which indicates photosynthesis and biomass production; the other is Secchi depth, a measure of water clarity or a measure of turbidity, reflecting planktonic growth in the absence of inorganic suspended material. In many marine and estuarine instances dissolved oxygen concentration (DO) and macrophyte growth and density are also important measures and, where indicated, may be included as initial response variables. Other measures can also be used, but these have been selected by EPA as of primary concern.

Nutrient criteria consist of judicious incorporation of present **reference condition information** about the primary variables, together with a **knowledge of historical conditions** and trends in the nutrient quality of the resource. These two factors, possibly augmented by **data extrapolations or models**, are analyzed objectively by a **panel of regional specialists** well versed in the biology, physics, and chemistry of the systems of concern. The criteria are also evaluated with respect to the **possible consequences of their implementation on downstream receiving waters**. All of these elements are required for the development of a nutrient criterion.

With this information, the status of a given water resource can be determined, management plans can be made, and management efforts can be evaluated.

The best possible understanding of the physical, chemical, and biological interrelationships in the environment is important in nutrient criteria development and the subsequent management response. However, effective nutrient criteria can and should be developed even in the absence of an in-depth scientific investigation of the ecological processing of nutrients in the estuarine and marine environment. An adequate number of proximal reference sites and current knowledge of the system are sufficient to initiate criteria development and proposed management responses. A conservative, environmentally responsible start can be made toward alleviating nutrient pollution, subject to adjustment as more scientific knowledge is obtained and verified.

The reference condition approach to criteria development was peer reviewed by the USEPA Science Advisory Board in 1990 and 1994 and judged to be scientifically defensible. It is also likely to be the most cost-effective approach.

EXECUTIVE SUMMARY

This manual is designed for use by State, Tribal, and Federal water resource managers as they address the cultural enrichment of their waters in conjunction with the EPA National Nutrient Criteria Program. It is intended to provide a stepwise sequence of actions leading to the development of nutrient criteria for estuarine and near-coastal marine waters to be used in correcting this overenrichment problem.

The premise of the National Nutrient Criteria Program is that many, if not most, of our nation's estuarine and coastal waters are moderately to severely polluted by excessive nutrients (Bricker et al.1999), especially nitrogen and phosphorus. This nutrient pollution affects not only the biotic integrity of the waters and the decline of valuable fish and shellfish, it has the potential to cause harm to the public health through hazardous algal blooms and the propagation of waterborne diseases. To address this problem, EPA uses a regionalized, waterbody type specific approach to the development of nutrient criteria or benchmarks for management decisionmaking. These criteria are based on the measurement of the most natural (or least impacted by human development) waters of a given type in a given area reflecting the condition to be expected in that region if human impacts are not a factor or are at least minimized. The variables of specific concern are total phosphorus and total nitrogen as causal variables, algal biomass (e.g., chlorophyll *a* for phytoplankton and ash-free dry weight for macroalgae), and water clarity (e.g., Secchi depth) as early response variables. In waters that already experience hypoxia, dissolved oxygen should be added as a response variable. EPA encourages States and Tribes to consider additional response indicators such as seagrasses and algal species composition.

This natural ambient background or "reference condition" is an important element of the nutrient criteria to be developed. The other elements are: an understanding of the historical status and trend of the water resource to help put the reference condition in perspective; models of the nutrient data to help better understand historical and present information and to project future consequences; concern and attention to the effects of any criteria development on downstream receiving waters; and the objective compilation and assessment of all of this information by a skilled body of regional experts...the "Regional Technical Assistance Group" or RTAG. The regional criteria so developed are guidelines the States and Tribes of the continental United States can use as they prepare their own criteria and standards for the improvement and protection of the nation's coastal waters.

The first of the actions needed to reach this criteria objective is the organization and utilization in each EPA Region of an RTAG consisting of specialists from State and Federal natural resource management agencies versed in the management and scientific principles most appropriate to that region and those waters. These are water resource managers, oceanographers, chemists, land use specialists, biologists, estuarine ecologists, statisticians, and similar local civil service experts employed by the State or Federal government. Academicians, special interest groups, and environmental group representatives are also important participants in the criteria development process and may assist the RTAG in its efforts.

The first requirement of the RTAG is the review and refinement of ecoregional determinations as most appropriate to the area. These are the geographic boundaries surrounding the similar estuarine and

coastal marine waters for which the criteria will apply. They are based on the EPA Ecoregion concept and incorporate attendant coastal Provinces, both of which are based on geographic and geologic similarities of landforms and parent material. The importance of this regionalization is the effort to deal with waters all having a similar inherent background nutrient loading and response characteristic. Once the regional boundaries and perhaps subregional divisions are completed, the RTAG investigates the physical classification of the waters into similar estuaries or coastal reaches or embayments for criteria development. In many instances the estuaries may be unique and require specific criteria.

Within the classification scheme developed, reference sites are identified as those areas suffering the least cultural development or impact, and the compilation of similar reference sites becomes a reference condition. The manual describes the scientific rationale for the variables selected, the dynamics of the receiving waters, and potentially confounding physical and chemical interrelationships influencing criteria development. It also describes sampling and analytical techniques for data gathering and processing to develop the reference conditions as well as several options for the compiling of this information. These include: (1) recognition and measurement of an excellent water body of ideal nutrient water quality with the aim of preserving this state; (2) in situ reference site determinations for moderately degraded waters; (3) hind casting for historical information from past higher nutrient quality conditions to determine the reference condition when no reference sites remain; (4) use of loading estimations from reference quality subestuarine tributary systems and projection to the estuary; and (5) options for establishing coastal nutrient reference conditions including a Nutrient Criteria Program pilot demonstration project.

Once the reference condition(s) has been determined, the RTAG then addresses the historical perspective; considers the need for models to project future consequences; considers the potential effect on receiving waters; and employs its own good judgment in collectively determining the appropriate criteria values for each of the variables to protect the waters of concern and their designated uses. A procedure is also suggested to equate the multiple criteria variables in a comprehensive dimensionless index score. The manual concludes with a chapter on model development and applications to the criteria program, and a chapter describing the application and implementation of nutrient criteria with emphasis on EPA Standards and Monitoring Divisions and a description of a comprehensive ten step sequential technique for water resource management.

This comprehensive progression from data collection to reference condition determination to criteria development and management responses, is intended to help users achieve the restoration and protection of the nutrient water quality of the nation's estuarine and near-coastal marine water resources.